

## NF-KB P65 RABBIT PAB

货号: N225275

产品全名: NF-KB p65 兔多抗

基因符号 NFKB3; RELA; TF65; Transcription factor p65; p65; NFkB

**UNIPROT ID:** Q04206

背景: NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL (MIM 164910), RELA, or RELB (MIM 604758) to form the NFKB complex. The p50 (NFKB1)/p65 (RELA) heterodimer is the most abundant form of NFKB. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, MIM 164008 or NFKBIB, MIM 604495), which inactivate NFKB by trapping it in the cytoplasm.

抗原: Peptide sequence around aa.309~313 (F-K-S-I-M) derived from Human NFkB-p65.

经过测试的应用: WB,IHC-P

推荐稀释比: WB: 1/500-1/1000 IHC: 1/50-1/100

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

分子量: Calculated MW: 60 kDa; Observed MW: 65 kDa

亚型: IgG


纯化: Affinity Chromatography

种属反应性: Human, Mouse and Rat


成分: PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

研究领域: Cell Biology

储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Western blot analysis of NF-KB p65 in HeLa, Raji and 3T3 lysates using NF-KB p65 antibody



Immunohistochemistry analysis of paraffin-embedded Human breast carcinoma tissue using NF-KB p65 antibody. Sample with blocking peptide on the right. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.