

PPAR DELTA (4F4) MOUSE MAB

货号: N261392

产品全名: PPAR delta (4F4) 小鼠单抗

基因符号 FAAR; NR1C2; NUC1; Peroxisome proliferative activated receptor delta

UNIPROT ID: Q03181

背景: Ligand-activated transcription factor. Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Has a preference for poly-unsaturated fatty acids, such as gamma-linoleic acid and eicosapentanoic acid. Once activated by a ligand, the receptor binds to promoter elements of target genes. Regulates the peroxisomal beta-oxidation pathway of fatty acids. Functions as transcription activator for the acyl-CoA oxidase gene. Decreases expression of NPC1L1 once activated by a ligand.

抗原: Purified recombinant protein expressed in E.coli.

经过测试的应用: IHC-P

推荐稀释比: IHC: 1/50-1/100

种属反应性: Mouse

克隆性: Mouse Monoclonal

克隆编号: 4F4-3B6-7D1

分子量: -

亚型: IgG1


纯化: Affinity Purified

种属反应性: Human,Rat,Mouse


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

研究领域: Epigenetics and Nuclear Signaling

储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemical analysis of paraffin-embedded Human tonsils using PPAR delta (4F4) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemistry analysis of paraffin-embedded Human Brain Tissue using PPAR delta (4F4) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.