

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

FLT3LG (DMC421) IGG1 CHIMERIC MAB

货号: 28215

产品全名: FLT3LG(DMC421) IgGI Chimeric 单抗

基因符号 FLT3LG; FL; FLT3L; Flt3 ligand

描述: FLT3LG antibody(DMC421) IgGI Chimeric 单抗

背景: FMS-like tyrosine kinase 3 ligand (Flt-3 Ligand) is also known as FL; Flt3L and FLT3LG; is an a-helical cytokine that promotes the differentiation of multiple hematopoietic cell lineages. FLT3LG is expressed as a noncovalentlylinked dimer by T cells and bone marrow and thymic fibroblasts. Each 36 kDa chain carries approximately 12 kDa of N- and Olinked carbohydrates. FLT3LG is structurally homologous to stem cell factor (SCF) and colony stimulating facor 1 (CSF-1). FLT3LG acts as a growth factor that increases the number of immune cells by activating the hematopoietic progenitors. It also induces the mobilization of the hematopoietic progenitors and stem cells in vivo which may help the system to kill cancer cells. FLT3LG induces the expansion of monocytes and immature dendritic cells as well as early B cell lineage differentiation. FLT3LG cooperates with IL2; IL6; IL7; and IL15 to induce NK cell development and with IL3; IL7 and IL11 to induce terminal B cell maturation. Animal studies also show FLT3LG to reduce the severity of experimentally induced allergic inflammation. FLT3LG is crucial for steady-state pDC and cDC development. A lack of FLT3L results in low levels of DCs.

经过测试的应用: Flow Cyt

推荐稀释比: Flow Cyt 1:100

种属反应性: Rabbit

亚型: Rabbit:Human Fc chimeric IgG1

纯化: Purified from cell culture supernatant by affinity chromatography 种属反应性: 人 FLT3LG

成分: Lyophilized from sterile PBS, pH 7.4. 5 % - 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).



Product Description

Pioneering GTPase and Oncogene Product Development since 2010




Figure 1. Flow cytometry analysis with Anti-FLT3LG (DMC421) on Expi293 cells transfected with human FLT3LG (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).