

## HUMAN CCL20 PROTEIN, HFC TAG

货号: 11238

产品全名: 人 CCL20 蛋白

规格: 10/50/100 µg

基因符号 CKb4;Exodus;LARC;MIP-3-alpha;MIP-3a;MIP3A;SCYA20;ST38

目标蛋白: CCL20

**UNIPROT ID:** P78556

**描述:** Recombinant Human CCL20 Protein with C-terminal human Fc tag

**背景:** This antimicrobial gene belongs to the subfamily of small cytokine CC genes. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The protein encoded by this gene displays chemotactic activity for lymphocytes and can repress proliferation of myeloid progenitors. Two transcript variants encoding different isoforms have been found for this gene.

**物种/宿主:** HEK293

**分子量:** The protein has a predicted molecular mass of 34.3 kDa after removal of the signal peptide. The apparent molecular mass of CCL20-hFc is approximately 35-55 kDa due to glycosylation.

**分子特征:** CCL20(Ala27-Met96) hFc(Glu99-Ala330)

**纯化:** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

**Formulation & Reconstitution:** Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization.

**储存和运输:** Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human CCL20 Protein, hFc Tag on SDS-PAGE under reducing condition.