HUMAN CD68 PROTEIN, HIS TAG

: 11397

: CD68

: 10/50/100 μg

GP110;LAMP4;SCARD1

: CD68

UNIPROT ID: P34810

: Recombinant human CD68 protein with C-terminal 6xHis tag

: This gene encodes a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It is a member of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family. The protein primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. It is a type I integral membrane protein with a heavily glycosylated extracellular domain and binds to tissue- and organ-specific lectins or selectins. The protein is also a member of the scavenger receptor family. Scavenger receptors typically function to clear cellular debris, promote phagocytosis, and mediate the recruitment and activation of macrophages. Alternative splicing results in multiple transcripts encoding different isoforms. [provided by RefSeq, Jul 2008]

/ : HEK293

: The protein has a predicted molecular mass of 32.4 kDa after removal of the signal peptide. The apparent molecular mass of CD68-His is approximately 55-130 kDa due to glycosylation.

: CD68(Asn22-Ser319) 6×His tag

: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

: Store at -20ºC to -80ºC for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80ºC (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Figure 1. Human CD68 Protein, His Tag on SDS-PAGE under reducing condition.