

HUMAN CXCL12 PROTEIN, HFC TAG

货号: 11679

产品全名: 人 CXCL12 蛋白

规格: 10/50/100 µg

基因符号 IRH;PBSF;SCYB12;SDF1;TLSF

目标蛋白: CXCL12

UNIPROT ID: P48061

描述: Recombinant Human CXCL12 with N-terminal human Fc tag

背景: This antimicrobial gene encodes a stromal cell-derived alpha chemokine member of the intercrine family. The encoded protein functions as the ligand for the G-protein coupled receptor, chemokine (C-X-C motif) receptor 4, and plays a role in many diverse cellular functions, including embryogenesis, immune surveillance, inflammation response, tissue homeostasis, and tumor growth and metastasis. Mutations in this gene are associated with resistance to human immunodeficiency virus type 1 infections. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2014]

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 34.7 kDa after removal of the signal peptide. The apparent molecular mass of hFc-CXCL12 is approximately 35-55 kDa due to glycosylation.

分子特征: hFc(Glu99-Ala330) CXCL12(Lys22-Met93)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human CXCL12 Protein, hFc Tag on SDS-PAGE under reducing condition.